

LETTER TO THE EDITOR

The Hamiltonian mass of asymptotically anti-de Sitter space-times

Piotr T Chruściel¹ and Gabriel Nagy²

Département de Mathématiques, Faculté des Sciences, Parc de Grandmont, F-37200 Tours, France

E-mail: chrusciel@univ-tours.fr and nagy@gargan.math.univ-tours.fr

Received 18 January 2001

Abstract

We give a Hamiltonian definition of mass for spacelike hypersurfaces in space-times with metrics which are asymptotic to the anti-de Sitter one, or to a class of generalizations thereof. We present the results of another paper (Chruściel P T and Nagy G 2001b, in preparation) which show that our definition provides a geometric invariant for a spacelike hypersurface \mathcal{S} embedded in a space-time (\mathcal{M}, g) . Some further global invariants are also given.

PACS numbers: 1130, 0420, 0240M

Let \mathcal{S} be an n -dimensional spacelike hypersurface in a $(n + 1)$ -dimensional Lorentzian space-time (\mathcal{M}, g) . Suppose that \mathcal{M} contains an open set \mathcal{U} which is covered by a finite number of coordinate charts (t, r, v^A) , with $r \in [R, \infty)$, and with (v^A) , local coordinates on some compact $(n - 1)$ -dimensional manifold M , such that $\mathcal{S} \cap \mathcal{U} = \{t = 0\}$. Assume that the metric g approaches a background metric b of the form

$$\begin{aligned} b &= -a^{-2}(r)dt^2 + a^2(r)dr^2 + r^2h, \\ h &= h_{AB}(v^C)dv^A dv^B, \end{aligned} \quad (1)$$

with $a(r) = 1/\sqrt{r^2/\ell^2 + k}$, where h is a Riemannian metric on M , k is a constant, and ℓ is a strictly positive constant related to the cosmological constant Λ by the formula $2\Lambda = -n(n - 1)/\ell^2$ (somewhat more general metrics are considered in [14]). For example, if h is the standard round metric on S^2 and $k = 1$, then b is the anti-de Sitter metric. To make the approach rates precise it is convenient to introduce an orthonormal frame for b ,

$$e_0 = a(r)\partial_t, \quad e_1 = \frac{1}{a(r)}\partial_r, \quad e_A = \frac{1}{r}\beta_A, \quad (2)$$

with β_A , an h -orthonormal frame on (M, h) , so that $b_{ab} = b(e_a, e_b) = \eta_{ab}$, the usual Minkowski matrix $\text{diag}(-1, +1, \dots, +1)$. We then require that the frame components g_{ab}

¹ Supported in part by the Polish Research Council grant KBN 2 P03B 073 15.

² Supported by a grant from Région Centre.

of g with respect to the frame (2) satisfy³

$$e^{ab} = O(r^{-\beta}), \quad e_a(e^{bc}) = O(r^{-\beta}), \quad b_{ab}e^{ab} = O(r^{-\gamma}), \quad (3)$$

where $e^{ab} = g^{ab} - b^{ab}$, with

$$\beta > n/2, \quad \gamma > n. \quad (4)$$

(The $(n + 1)$ dimensional Schwarzschild–anti de Sitter metrics (15) satisfy (3) with $\beta = n$, and with $\gamma = 2n$. The boundary conditions (3)-(4), and the overall approach here stem from [15, section 5].) We stress that we do *not* assume the existence of global frames on the asymptotic region: when M is not parallelizable, then conditions (3)-(4) should be understood as the requirement of *the existence of a covering of M by a finite number of open sets \mathcal{O}_i together with frames defined on $[R_0, \infty) \times \mathcal{O}_i$ satisfying (3)-(4)*. For simplicity, further suppose that g satisfies the vacuum Einstein equations with a cosmological constant,

$$R_{\mu\nu} - \frac{g^{\alpha\beta}R_{\alpha\beta}}{2}g_{\mu\nu} = -\Lambda g_{\mu\nu}, \quad (5)$$

similarly for b . (The existence of a large family of such g 's follows from the work in [17, 24].) A geometric Hamiltonian analysis [10, 14], based on the formalism of [26], leads to the following expression for the Hamiltonian associated to the flow of a vector field X , assumed to be a Killing vector field of the background b :⁴

$$m(\mathcal{S}, g, b, X) = \frac{1}{2} \int_{\partial\mathcal{S}} \mathbb{U}^{\alpha\beta} dS_{\alpha\beta}, \quad (6)$$

$$\mathbb{U}^{\nu\lambda} = \mathbb{U}^{\nu\lambda}{}_{\beta} X^{\beta} + \frac{1}{8\pi} \left(\sqrt{|\det g_{\rho\sigma}|} g^{\alpha[\nu} - \sqrt{|\det b_{\rho\sigma}|} b^{\alpha[\nu} \right) \delta_{\beta}^{\lambda]} \overset{\circ}{\nabla}_{\alpha} X^{\beta}, \quad (7)$$

$$\mathbb{U}^{\nu\lambda}{}_{\beta} = \frac{2|\det b_{\mu\nu}|}{16\pi \sqrt{|\det g_{\rho\sigma}|}} g_{\beta\gamma} \overset{\circ}{\nabla}_{\kappa} (e^2 g^{\gamma[\nu} g^{\lambda]\kappa}), \quad (8)$$

$$e = \sqrt{|\det g_{\rho\sigma}|} / \sqrt{|\det b_{\mu\nu}|}. \quad (9)$$

It is appropriate to make a few comments: in [10] the starting point of the analysis is the Hilbert Lagrangian for vacuum Einstein gravity, $\mathcal{L} = \sqrt{-\det g_{\mu\nu}} g^{\alpha\beta} R_{\alpha\beta} / 16\pi$. With our signature $(- + \dots +)$ one needs to repeat the analysis in [10] with \mathcal{L} replaced by $\sqrt{-\det g_{\mu\nu}} (g^{\alpha\beta} R_{\alpha\beta} - 2\Lambda) / 16\pi$, and without making the assumption $n + 1 = 4$. The final expression for the Hamiltonian (6) coincides with that obtained when $\Lambda = 0$ and $n + 1 = 4$. Equation (7) coincides with that derived in [10] except for the term $-\sqrt{|\det b_{\rho\sigma}|} b^{\alpha[\nu} \delta_{\beta}^{\lambda]} \overset{\circ}{\nabla}_{\alpha} X^{\beta}$. This term does not depend on the metric g , and such terms can be freely added to the Hamiltonian because they do not affect the variational formula that defines a Hamiltonian. From an energy point of view such an addition corresponds to a choice of the zero point of the energy. We note that in our context $m(\mathcal{S}, g, b, X)$ would not converge if the term $-\sqrt{|\det b_{\rho\sigma}|} b^{\alpha[\nu} \delta_{\beta}^{\lambda]} \overset{\circ}{\nabla}_{\alpha} X^{\beta}$ was not present in (7). The extension to non-vacuum models is

³ The summation convention is used throughout. We use Greek indices for coordinate components and lower-case Latin indices for the tetrad ones; upper-case Latin indices run from 2 to n and are associated either to coordinates or to frames on M , in a way which should be obvious from the context. Finally Greek *bracketed* indices (α) etc refer to objects defined outside of space-time, such as the Killing algebra $\mathcal{K}_{\mathcal{S}^{\perp}}$, or an exterior embedding space, and are also summed over.

⁴ The integral over $\partial\mathcal{S}$ should be understood by a limiting process, as the limit as R tends to infinity of integrals over the sets $t = 0, r = R$. $dS_{\alpha\beta}$ is defined as $\frac{\partial}{\partial x^{\alpha}} \rfloor \frac{\partial}{\partial x^{\beta}} \rfloor dx^0 \wedge \dots \wedge dx^n$, with \rfloor denoting contraction; g stands for the space-time metric unless explicitly indicated otherwise. Square brackets denote antisymmetrization with an appropriate numerical factor (1/2 for two indices), and $\overset{\circ}{\nabla}$ denotes covariant differentiation *with respect to the background metric b* .

straightforward and leads to the same Hamiltonian for several standard matter models; the simplest way to prove this is by following the arguments in [25].

We show in [14] that the vacuum Einstein equations together with the decay conditions (3)-(4) and the asymptotic behaviour of the frame components of the b -Killing vector fields,

$$X^a = O(r), \quad \hat{\nabla}_a X^b = O(r),$$

guarantee that the integrals (6) are convergent. Further, it is easy to see that under the asymptotic conditions (3)-(4) the volume integrals appearing in the variational formulae in [10] are convergent, the undesirable boundary integrals in the variational formulae of [10] vanish, so that the integrals (6) do indeed provide Hamiltonians on the space of fields satisfying (3)-(4). (Compare with [20] for an alternative, essentially equivalent, Hamiltonian approach.) This singles out the charges (6) amongst various alternative expressions because Hamiltonians are uniquely defined, up to the addition of a constant, on each path connected component of the phase space. The key advantage of the Hamiltonian approach is precisely this uniqueness property, which does not seem to have a counterpart in the Noether charge analysis [29] (cf, however [30, 23]), or in Hamilton–Jacobi type arguments [8].

To define the integrals (6) we have fixed a model background metric b , as well as an orthonormal frame as in (2). This last equation requires the corresponding coordinate system (t, r, v^A) as in (1). Hence, the background structure required in our analysis consists of a *background metric* and a *background coordinate system*. While this is reasonably satisfactory from a Hamiltonian point of view, in which each choice of background structure defines an associated phase space, there is an essential *potential geometric ambiguity* in the integrals (6) that arises as follows: let g be any metric such that its frame components g^{ab} tend to η^{ab} as r tends to infinity, in such a way that the integrals $m(\mathcal{S}, g, b, X)$ given by (6) (labelled by all the background Killing vector fields X or perhaps by a subset thereof) converge. Consider another coordinate system $(\hat{t}, \hat{r}, \hat{v}^A)$ with the associated background metric \hat{b} :

$$\hat{b} = -a^{-2}(\hat{r})d\hat{t}^2 + a^2(\hat{r})d\hat{r}^2 + \hat{r}^2\hat{h}, \quad \hat{h} = h_{AB}(\hat{v}^C)d\hat{v}^A d\hat{v}^B,$$

together with an associated frame \hat{e}^a ,

$$\hat{e}_0 = a(\hat{r})\partial_{\hat{t}}, \quad \hat{e}_1 = \frac{1}{a(\hat{r})}\partial_{\hat{r}}, \quad \hat{e}_A = \frac{1}{\hat{r}}\hat{\beta}_A, \quad (10)$$

and suppose that in the new hatted coordinates the integrals defining the charges $m(\hat{\mathcal{S}}, g, \hat{b}, \hat{X})$ converge again. An obvious way of obtaining such coordinate systems is to make a coordinate transformation

$$t \rightarrow \hat{t} = t + \delta t, \quad r \rightarrow \hat{r} = r + \delta r, \quad v^A \rightarrow \hat{v}^A = v^A + \delta v^A, \quad (11)$$

with $(\delta t, \delta r, \delta v^A)$ decaying sufficiently fast:

$$\begin{aligned} \hat{t} &= t + O(r^{-1-\beta}), & e_a(\hat{t}) &= \ell \delta_a^0 + O(r^{-1-\beta}), \\ \hat{r} &= r + O(r^{1-\beta}), & e_a(\hat{r}) &= \frac{\delta_a^1}{\ell} + O(r^{1-\beta}), \\ \hat{v}^A &= v^A + O(r^{-1-\beta}), & e_a(\hat{v}^A) &= \delta_a^A + O(r^{-1-\beta}), \end{aligned} \quad (12)$$

and with analogous conditions on second derivatives; this guarantees that the hatted analogue of equations (3) and (4) will also hold. In [14] we prove that under coordinate transformations (12) the integrals (6) remain unchanged:

$$m(\mathcal{S}, g, b, X) = m(\hat{\mathcal{S}}, g, \hat{b}, \hat{X}).$$

Here, if $X = X^\mu(t, r, v^A)\partial_\mu$, then the vector field \hat{X} is defined using the *same* functions X^μ of the *hatted* variables. The proof is a computation of the change of the integrand of (6)

under the change of coordinates (11). One notices that all the terms obtained by linearizing in $(\delta t, \delta r, \delta v^A)$ can be written as a total divergence, and therefore give no contribution; this is reminiscent of a divergence identity which is used for a similar calculation for asymptotically flat space-times. One finally checks that under the boundary conditions (3)-(4) the correction terms give vanishing contribution in the limit $r \rightarrow \infty$.

This result appears to be sharp: indeed, consider the metric $g = \hat{b}$, with the hatted coordinates defined as

$$\hat{r} = r + \frac{\zeta}{r^{n/2-1}}, \quad \hat{v}^A = v^A, \quad (13)$$

where ζ is a constant. Thus, if \hat{b} is the anti-de Sitter metric, then g is again the anti-de Sitter metric in a different coordinate system, which does not differ too much from the original one, in particular the metric approaches asymptotically the standard version of the anti-de Sitter metric in the unhatted coordinates. Then the components of g with respect to the ON frame associated to the unhatted background b satisfy (3) with $\beta = \gamma = n/2$. A calculation shows that g has a non-vanishing mass integral (6) with respect to the unhatted background b . The same coordinate transformation leads to a non-zero Abbott–Deser [1] mass of $g = \hat{b}$ with respect to b .

It should be stressed that *a priori* we did not know that the hatted coordinates were related to the unhatted ones by the simple coordinate transformation (11) with $(\delta t, \delta r, \delta v^A)$ decaying as $r \rightarrow \infty$, or that they behaved in some controlled way, indeed the behaviour of $(\delta t, \delta r, \delta v^A)$ could in principle be very wild. The main technical result of [14] is the proof that this is not the case: under the hypothesis that (M, h) is a sphere with a round metric, or that (M, h) has a non-positive Ricci tensor and constant Ricci scalar⁵ we show that *all coordinate transformations which leave \mathcal{S} invariant (so that $\hat{t} = t$) and that preserve the decay conditions (3)-(4) are compositions of a map satisfying (12) with an isometry of the background*. In order to obtain a geometric invariant of \mathcal{S} it thus remains to study the behaviour of the integrals (6) under isometries of b preserving \mathcal{S} . If Φ is such an isometry, the fact that $\mathbb{U}^{\alpha\beta}$ is a tensor density immediately yields the formula

$$m(\mathcal{S}, \Phi^*g, b, (\Phi_*)^{-1}X) = m(\mathcal{S}, g, b, X). \quad (14)$$

(Here Φ^* is the pull-back, and Φ_* is the push-forward map. When understood in a passive manner, Φ^*g is simply the metric g expressed in the new coordinates, while $(\Phi_*)^{-1}X$ is the vector field X expressed in the new coordinates.) This has the effect that the *integrals (6) are reshuffled amongst each other under isometries of b , in a way determined by the representation of the group of isometries of b on the space of Killing vectors*, and leads to the following:

- (a) Let M be the $(n - 1)$ -dimensional sphere ${}^{n-1}S$ with a round metric h , normalized so that the substitution $k = 1$ in (1) leads to a metric b which is the $(n + 1)$ -dimensional anti-de Sitter metric. The space $\mathcal{K}_{\mathcal{S}^\perp}$ of b -Killing vector fields normal to $\mathcal{S} \cap \mathcal{U}$ is spanned by vector fields $K_{(\mu)}$ which on \mathcal{S} take the form $K_{(\mu)} = N_{(\mu)}e_0$, $\mu = 0, \dots, n$, where $N_{(0)} = \sqrt{\frac{r^2}{\ell^2} + 1}$, $N_{(i)} = \frac{x^i}{\ell}$, and $x^i = rn^i$, r being the coordinate which appears in (1), while $n^i \in {}^{n-1}S \subset \mathbb{R}^n$. The group $\text{Iso}(\mathcal{S}, b)$ of isometries Φ of b which map \mathcal{S} into \mathcal{S} coincides with the homogeneous Lorentz group $O(1, n)$; it acts on $\mathcal{K}_{\mathcal{S}^\perp}$ by push-forward. It can be shown that for each such Φ there exists a Lorentz transformation $\Lambda = (\Lambda^{(\alpha)}_{(\beta)}) : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ so that for every $X = X^{(\alpha)}K_{(\alpha)} \in \mathcal{K}_{\mathcal{S}^\perp}$ we have

$$(\Phi_*X)^{(\alpha)} = \Lambda^{(\alpha)}_{(\beta)}X^{(\beta)}.$$

⁵ Related results under less restrictive conditions can be found in [14].

We set

$$m_{(\mu)} \equiv m(\mathcal{S}, g, b, K_{(\mu)}),$$

it follows that the number

$$m^2(\mathcal{S}, g) = |\eta^{(\mu)(\nu)} m_{(\mu)} m_{(\nu)}|,$$

where $\eta^{(\mu)(\nu)} = \text{diag}(-1, +1, \dots, +1)$ is the Minkowski metric on \mathbb{R}^{n+1} , is an invariant of the action of $\text{Iso}(\mathcal{S}, b)$. Further, if we define $m(\mathcal{S}, g)$ to be positive if $m^{(\mu)} := \eta^{(\mu)(\nu)} m_{(\nu)}$ is spacelike, and we take the sign of $m(\mathcal{S}, g)$ to coincide with that of $m^{(0)}$ if $m^{(\mu)}$ is timelike or null, then $m(\mathcal{S}, g)$ is invariant under the action of the group $\text{Iso}^\uparrow(\mathcal{S}, b)$ of time-orientation preserving Lorentz transformations. The number $m(\{t = 0\}, g)$ so defined coincides with the mass parameter m of the Kottler ('Schwarzschild-anti-de Sitter') metrics in dimension $(n+1) = 4$, and it is proportional to the parameter m which occurs in the $(n+1)$ -dimensional generalizations of the Kottler metrics [22]

$$g = - \left(1 - \frac{2m}{r^{n-2}} + \frac{r^2}{\ell^2} \right) dt^2 + \left(1 - \frac{2m}{r^{n-2}} + \frac{r^2}{\ell^2} \right)^{-1} dr^2 + r^2 h, \quad (15)$$

with h , a round metric of scalar curvature $(n-1)(n-2)$ on a $(n-1)$ -dimensional sphere (cf, e.g., [9]).

Consider the remaining Killing vector fields $L_{(\alpha)(\beta)}$ of b :

$$L_{(\alpha)(\beta)} = \eta_{(\alpha)(\sigma)} y^{(\sigma)} \frac{\partial}{\partial y^{(\beta)}} - \eta_{(\beta)(\sigma)} y^{(\sigma)} \frac{\partial}{\partial y^{(\alpha)}}. \quad (16)$$

Here the coordinates $y^{(\alpha)}$ are the coordinates on \mathbb{R}^{n+1} obtained by isometrically embedding, into \mathbb{R}^{n+1} , the $\{t = 0\}$ section of $(n+1)$ -dimensional anti-de Sitter space-time equipped with the Minkowski metric $\eta^{(\mu)(\nu)}$. It can be checked that under the action of $\text{Iso}(\mathcal{S}, b)$ the integrals

$$Q_{(\mu)(\nu)} \equiv m(\mathcal{S}, g, b, L_{(\mu)(\nu)})$$

transform as the components of a two-covariant antisymmetric tensor. One then obtains a geometric invariant of \mathcal{S} by calculating

$$Q \equiv Q_{(\mu)(\nu)} Q_{(\alpha)(\beta)} \eta^{(\mu)(\alpha)} \eta^{(\nu)(\beta)}. \quad (17)$$

In dimension $(3+1)$ another independent global geometric invariant is obtained from

$$Q^* \equiv Q_{(\mu)(\nu)} Q_{(\alpha)(\beta)} \epsilon^{(\mu)(\alpha)(\nu)(\beta)}. \quad (18)$$

In higher dimensions further invariants are obtained by calculating $\text{tr}(P^{2k})$, $2 \leq 2k \leq (n+1)$, where $P^{(\alpha)(\beta)} = \eta^{(\alpha)(\mu)} Q_{(\mu)(\beta)}$. (In this notation Q given by equation (17) equals $\text{tr}(P^2)$.)

- (b) Let M be a compact $(n-1)$ -dimensional manifold with a metric h of constant scalar curvature and with a non-positive Ricci tensor, and let b take the form (1), with $a(r) = 1/\sqrt{r^2/\ell^2 + k}$, and with $k = 0$ or -1 according to whether the Ricci scalar of h vanishes or not. We show in [14] that for such metrics the space of b -Killing vector fields normal to \mathcal{S} consists of vector fields of the form $X(\lambda) = \lambda \partial_t$, $\lambda \in \mathbb{R}$, and that

$$m(\mathcal{S}, g) \equiv m(\mathcal{S}, g, b, X(1))$$

is background independent, hence a geometric invariant⁶. Some other geometric invariants can be obtained from the integrals (6) when Killing vectors which are not necessarily

⁶ If the scalar curvature of h vanishes, then Einstein equations require the constant k in the metric to vanish. In that case there arises an ambiguity in the definition of mass related to the possibility of rescaling t and r without changing the form of the metric, which rescales the mass. (This freedom does not occur when k is non-zero.) This ambiguity can be removed by arbitrarily choosing some normalization for the h -volume of M , e.g. 4π in dimension $(n+1) = 4$.

normal to \mathcal{S} exist, using invariants of the action of the isometry group of b on the space of Killing vectors. If the Ricci tensor of M is strictly negative no other Killing vectors exist. On the other hand, if h is a flat torus, then each h -Killing vector provides a global geometric invariant via the integrals (6).

The number $m(\mathcal{S}, g)$ defined in each case above is our proposal for the geometric definition of total mass of \mathcal{S} in (\mathcal{M}, g) . We note that its multiplicative normalization in $(n + 1) = 4$ dimensions is determined by the requirement of the correct Newtonian limit when $M = S^2$ and $\Lambda = 0$, while the additive one is determined by imposing that the background models have vanishing energy. In higher dimensions it appears appropriate to keep the same multiplicative factor for the Lagrangian \mathcal{L} , whenever a Kaluza–Klein reduction applies.

The results described above can be reformulated in a purely Riemannian context, this will be discussed elsewhere [13]. Some similar results in that context have been obtained independently by M Herzlich [21].

It is natural to study the invariance of the mass when \mathcal{S} is allowed to move in \mathcal{M} . A complete answer would require establishing an equivalent of our analysis of admissible coordinate transformations in a space-time setting. The difficulties that arise in the corresponding problem for asymptotically Minkowskian metrics [12] suggest that this might be a considerably more delicate problem, which we plan to analyse in the future. It should be stressed that this problem mixes two different issues, one being the potential background dependence of (6), another one being the possibility of energy flowing in or out through the timelike conformal boundary of space-time.

It should be pointed out that there exists several alternative methods of defining mass in asymptotically anti-de Sitter space-time, using coordinate systems [7, 19], preferred foliations [18], generalized Komar integrals [27], conformal techniques [2, 3, 5], or *ad-hoc* methods [1]; an extended discussion can be found in [15, section 5]. Each of those approaches suffers from some *potential* ambiguities, so that the question of the geometric character of the definition of mass given there arises as well. Let us briefly compare the results described here to some of the work of others. Consider first, the Abbott–Deser approach [1], which seems to have been used most often. A precise comparison is difficult to carry out because in [1] the boundary conditions which should be used are not described in detail. As already pointed out, the coordinate transformation (13) gives a non-zero Abbott–Deser mass to the anti-de Sitter metric, so the boundary conditions *do* matter. Assuming that the authors of [1] had in mind boundary conditions which are at least as restrictive as our conditions (3)-(4), a straightforward but tedious calculation shows that the Abbott–Deser mass coincides with the Hamiltonian mass advocated here. One then needs to face the same ambiguities as we do, and our results in [14] can be interpreted as proving the existence of a geometric invariant which can be calculated using Abbott–Deser type integrals. It should be stressed, however, that while the Hamiltonian approach is universal and leads to unique functionals (up to a constant) on each phase space, no results about either the uniqueness or universality of the approach of [1] are known to us.

Consider next the Hamiltonian approach of [20]. Here the question of equivalence of the different Hamiltonian formalisms of [10] and of [20] arises. Those formalisms are identical in spirit, as both are Hamiltonian, but differ in various details. We note that the boundary conditions (3)-(4) are weaker than the conditions imposed in [20]. It can be checked that under (3)-(4) the formalism of [20] is a special case of the geometric Hamiltonian formalism of [10], when applied to asymptotically anti-de Sitter space-times, so the Hamiltonian part of our analysis in [14] can be thought of as an extension of the results of [20] to larger phase spaces arising naturally in this context. An advantage of the geometric Hamiltonian formalism of Kijowski and Tulczyjew is that it allows the use of both a manifestly four-dimensional

covariant formalism, and of a (3 + 1) ADM one. The potential ambiguities in a geometric definition of mass that occur in [20] are identical to the ones described above.

As another example we note the potential ambiguity in the mass defined by the conformal methods in [2, 3], related to the possibility of existence of smooth conformal completions which are *not* smoothly conformally equivalent. In those works one assumes existence of smooth completions, while our conditions would, roughly speaking, correspond to $C^{1,\alpha}$ conformal completions, $\alpha > 1/2$, hence our conditions (3)–(4) are satisfied in the setup of [2, 3] (compare with [15] for a detailed discussion in the static case). It can be shown [4] that the mass of [2, 3] coincides with the Abbott–Deser one and what has been said above implies that, under the asymptotic conditions of [2, 3], also coincides with the Hamiltonian mass described here. The results proved in [14] can be used to show that no inequivalent conformal completions of the kind considered in [2, 3] exist, establishing the invariant character of the framework of [2, 3].

Let us finally turn our attention to the AdS/CFT motivated definitions of mass (cf, e.g., [2, 5, 16, 28] and references therein). It seems that there might be a belief in the string community that, at least in odd space-time dimensions, there is no systematic way to start from the Einstein–Hilbert action and arrive unambiguously at conserved quantities. Our results in [14] show that such a belief is incorrect, in the space of metrics asymptotic to the backgrounds considered here, as well as for those which asymptote to the more general backgrounds considered in [14]. It should be pointed out, however, that the Hamiltonian approach gives Hamiltonians which are only defined up to a constant, which leaves ample room for non-zero ‘Casimir energies’, the occurrence of which might well be justified by other physical considerations. On the other hand, in a Hamiltonian approach the only natural choice of the additive constant seems to be zero. In any case, it is not clear whether or not the AdS/CFT based approaches lead to geometric invariants of hypersurfaces in space-times, in the sense presented here.

We note that a similar problem for the ADM mass of asymptotically flat initial data sets has been settled in [6, 11] (see also [12]). Our treatment in [14] is a non-trivial extension to the current setup of the methods of [11].

Acknowledgments

PTC wishes to thank M Herzlich, S Ilias, A Polombo and A El Soufi for useful discussions. Suggestions from an anonymous referee for improvements of the manuscript are also acknowledged.

References

- [1] Abbott L F and Deser S 1982 Stability of gravity with a cosmological constant *Nucl. Phys. B* **195** 76–96
- [2] Ashtekar A and Das S 2000 Asymptotically anti-de Sitter space-times: conserved quantities *Class. Quantum Grav.* **17** L17–L30
(Ashtekar A and Das S 1999 *Preprint* hep-th/9911230)
- [3] Ashtekar A and Magnon A 1984 Asymptotically anti-de Sitter space-times *Class. Quantum Grav.* **1** L39–L44
- [4] Ashtekar A 2000 Private communication
- [5] Balasubramanian V and Kraus P 1999 A stress tensor for anti-de Sitter gravity *Commun. Math. Phys.* **208** 413–28
- [6] Bartnik R 1986 The mass of an asymptotically flat manifold *Comm. Pure Appl. Math.* **39** 661–693
- [7] Boucher W, Gibbons G W and Horowitz G T 1984 Uniqueness theorem for anti-de Sitter spacetime *Phys. Rev. D* **30** 2447–2451
- [8] Brown J D and York J W Jr 1993 Quasilocal energy and conserved charges derived from the gravitational action *Phys. Rev. D* **47** 1407–19
- [9] Cadeau C and Woolgar E 2001 New five dimensional black holes classified by horizon geometry, and a Bianchi VI braneworld *Class. Quantum Grav.* **18** 527–42

- (Cadeau C and Woolgar E 2000 *Preprint* gr-qc/0011029)
- [10] Chruściel P T 1985 On the relation between the Einstein and the Komar expressions for the energy of the gravitational field *Ann. Inst. H. Poincaré* **42** 267–82
- [11] Chruściel P T 1986 Boundary conditions at spatial infinity from a hamiltonian point of view *Topological Properties and Global Structure of Space-Time* ed P Bergmann and V de Sabbata (New York: Plenum) pp 49–59 <http://www.phys.univ-tours.fr/~piotr/scans>
- [12] Chruściel P T 1988 On the invariant mass conjecture in general relativity *Commun. Math. Phys.* **120** 233–48
- [13] Chruściel P T and Nagy G 2001a The mass of spacelike hypersurfaces in asymptotically anti-de Sitter space-times, in preparation
- [14] Chruściel P T and Nagy G 2001b The mass of asymptotically hyperboloidal Riemannian manifolds, in preparation
- [15] Chruściel P T and Simon W 2001 Towards the classification of static vacuum spacetimes with negative cosmological constant *J. Math. Phys.* **42** 1779–817
(Chruściel P T and Simon W 2000 *Preprint* gr-qc/0004032)
- [16] Emparan R, Johnson C V and Myers R C 1999 Surface terms as counterterms in the AdS/CFT correspondence *Phys. Rev. D* **60** 104001
(Emparan R, Johnson C V and Myers R C 1999 *Preprint* hep-th/9903238)
- [17] Friedrich H 1995 Einstein equations and conformal structure: Existence of anti-de-Sitter-type space-times *J. Geom. and Phys.* **17** 125–184
- [18] Gibbons G W 1999 Gravitational entropy and the inverse mean curvature flow *Class. Quantum Grav.* **16** 1677–87
- [19] Gibbons G W, Hawking S W, Horowitz G T and Perry M J 1983 Positive mass theorems for black holes *Commun. Math. Phys.* **88** 295–308
- [20] Henneaux M and Teitelboim C 1985 Asymptotically anti-de Sitter spaces *Commun. Math. Phys.* **98** 391–424
- [21] Herzlich M 2000 Private communication
- [22] Horowitz G T and Myers R C 1999 The AdS/CFT correspondence and a new positive energy conjecture for general relativity *Phys. Rev. D* **59** 026005
- [23] Julia B and Silva S 2000 Currents and superpotentials in classical gauge theories. II: Global aspects and the example of affine gravity *Class. Quantum Grav.* **17** 4733–44
- [24] Kánnár J 1996 Hyperboloidal initial data for the vacuum Einstein equations with cosmological constant *Class. Quantum Grav.* **13** 3075–84
- [25] Kijowski J 1978 On a new variational principle in general relativity and the energy of the gravitational field *Gen. Rel. Grav.* **9** 857–77
- [26] Kijowski J and Tulczyjew W M 1979 A symplectic framework for field theories *Lecture Notes in Physics* **107** (New York: Springer)
- [27] Magnon A 1985 On Komar integrals in asymptotically anti-de Sitter space-times *J. Math. Phys.* **26** 3112–7
- [28] Myers R C 1999 Stress tensors and casimir energies in the AdS / CFT correspondence *Phys. Rev. D* **60** 046002
(Myers R C 1999 *Preprint* hep-th/9903203)
- [29] Trautman A 1962 Conservation laws in general relativity *Gravitation: An Introduction to Current Research* ed L Witten (New York: Wiley)
- [30] Wald R M and Zoupas A 2000 A general definition of ‘conserved quantities’ in general relativity and other theories of gravity *Phys. Rev. D* **61** 084027
(Wald R M and Zoupas A 1999 *Preprint* gr-qc/9911095)